CLH 302
Pollutant Contaminated

DATA REPORT

# EVALUATION OF TRACE-METAL LEVELS IN AMBIENT WATERS AND TRIBUTARIES TO NEW YORK/NEW JERSEY HARBOR FOR WASTE LOAD ALLOCATION

EPA Contract No. 68-C8-0105
Work Assignment No. 3-213

to

ENVIRONMENTAL PROTECTION AGENCY
Office of Wetlands, Oceans, and Watersheds
and
Region II

January 9, 1992

Prepared by
Battelle Ocean Sciences
397 Washington Street
Duxbury, MA 02332
(617) 934-0571

Newark Bay Study Area
NBSA-00071

CONTENTS


1.0 INTRODUCTION ..... 1

2.0 METHODS ..... 2
2.1 Sample Collection Procedures ..... 2
2.1.1 Ambient Water Samples ..... 2
2.2.2 Tributary Water Samples ..... 2
2.2 Sample Preparation ..... 2
2.2.1 Ambient Water Samples ..... 3
2.2.2 Tributary Water Samples ..... 4
2.2.3 Sample Digestion and Preconcentration ..... 5
2.3 Instrumental Analysis ..... 5
2.4 Sample Custody ..... 6
2.5 Data Quality Objectives ..... 6

3.0 RESULTS ..... 7

4.0 CONCLUSIONS ..... 8

5.0 REFERENCES ..... 9

Appendix - FIGURES AND TABLES

## LIST OF FIGURES


Figure 1. Station Locations for May 1991 Intensive Master Station Survey ..... A-1

Figure 2. (a) Total Suspended Solids in New York/New Jersey Harbor Waters ..... A-2
(b) Particulate Carbon and Dissolved Organic Carbon In New York/ New Jersey Harbor Waters ..... A-3
(c) Arsenic Concentrations in New York/New Jersey Harbor Waters ..... A-4
(d) Cadmium Concentrations in New York/New Jersey Harbor Waters ..... A-5
(e) Copper Concentrations in New York/New Jersey Harbor Waters ..... A-6
(f) Lead Concentrations in New York/New Jersey Harbor Waters ..... A-7
(g) Mercury Concentrations in New York/New Jersey Harbor Waters ..... A-8
(h) Nickel Concentrations in New York/New Jersey Harbor Waters ..... A-9
(i) Silver Concentrations in New York/New Jersey Harbor Waters ..... A-10
(j) Zinc Concentrations in New York/New Jersey Harbor Waters ..... A-11


## LIST OF TABLES


Table 1. List of samples collected during the May 1991 Intensive Master Station Survey ..... A-12

Table 2. Data-Quality Objective for Analytical Measurements ..... A-16

(a) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Salinity ..... A-17

(b) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Total Suspended Solids ..... A-18

(c) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Particulate Carbon and Dissolved Organic Carbon ... A-20

(d) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Arsenic ..... A-22

(e) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Cadmium ..... A-24

(f) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Copper ..... A-26

(g) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Lead ..... A-28

(h) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Mercury ..... A-30

(i) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Nickel ..... A-32

j) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Silver ..... A-34

(k) Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Zinc ..... A-36

Table 4. Precision Results of the Field-duplicate Samples Collected for the New York/New Jersey Harbor Intensive Master Station Survey — Relative Percent Difference A-38

Table 5. Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Surface Floc ..... A-39

# 1.0 INTRODUCTION

The Environmental Protection Agency (EPA) Region II and the States of New York and New Jersey have declared that the waters of New York/New Jersey Harbor (NY/NJ Harbor) are impaired under the provisions of Section 304(l) of the Federal Water Pollution Control Act. This determination was based on exceedances of EPA water quality criteria (WQC) and State standards for the metals copper and mercury. These WQC exceedances were attributed to point-source discharges, prompting the development of a pollution-control strategy that included an Intensive Master Station Survey (IMSS) to determine the concentrations of metals over a tidal cycle in surface and bottom layers of the water column and in the ambient waters of NY/NJ Harbor, and its major tributaries. The results of this survey are presented and briefly discussed in this report.

The objectives of this study were to provide data that could be used by EPA and others to

- Determine the tidal distribution and identify any patterns of metal concentrations in NY/NJ Harbor
- Determine the metals that exceed Federal or State water-quality criteria or standards and the locations of exceedances within NY/NJ Harbor
- Estimate the spatial distribution and identify any patterns of metal concentrations in NY/NJ Harbor
- Develop a water-quality based waste-load allocation model as part of the point-source discharge control strategy
- Determine metal concentration in particulate sample collected from approximately one meter above the sediment (surface floc).

To meet these objectives, a field survey was conducted in May 1991 to collect ambient water samples at six stations in the NY/NJ Harbor over a tidal cycle (see Figure 1 and Table 1. *Note*: All figures and tables are in the Appendix) and tributary samples at three New Jersey Tributaries: Hackensack, Passaic, and Raritan rivers (see Table 2). All samples were analyzed for trace metals (Ag, As, Cd, Cu, Hg, Ni, Pb, Zn), total suspended solids (TSS), particulate carbon (PC), dissolved organic carbon (DOC), and salinity. Metal concentrations were determined in two phases, total recoverable and dissolved, for all metals. In addition, mercury was analyzed in the particulate phase.

Sample collection was a joint effort among personnel from the New Jersey Department of Environmental Protection and Engineering (NJDEP&E), EPA, and Battelle.

Additional background information on this project is given in the Work/Quality Assurance Project Plans (W/QAPjPs; Battelle, 1990a) and Survey Plan (Battelle, 1990b). Activities that were conducted during the IMSS are summarized in a Survey Report Letter (Battelle, 1991c).

# 2.0 METHODS

## 2.1 SAMPLE COLLECTION PROCEDURES

### 2.1.1 Ambient Water Samples

All ambient water samples were collected by using the Battelle Ocean Sampling System (BOSS) on board the OSV *Anderson*. Details of the sample-collection procedures, including decontamination of the BOSS tubing, have been given in the Survey Plan (Battelle, 1990b). Sample aliquots were placed in polyethylene bottles for trace metals (two bottles), PC/DOC, and TSS measurements. Two Teflon® bottles were filled for Hg analysis. The bottles used for Hg and the other trace metals were cleaned by using the multistep process described by Patterson and Settle (1976). In addition, *in situ* measurements of (1) salinity, (2) temperature, (3) depth, and (4) transmissometry were obtained at each sampling event. Only salinity and depth are presented in this report.

### 2.1.2 Tributary Water Samples

All of the sample bottles required for a tributary were placed capless in an alconox-washed, deionized water-rinsed weighted, polypropylene-mesh bag. The bottles and mesh bags for each tributary were stored in clean plastic bags prior to and after use. For sampling, each mesh bag was secured with nylon rope and lowered down into the water. All bottle caps were kept in a clean plastic bag while the sample was being collected. The weighted bag was heavy enough to submerge all bottles at the same time to a depth of about 1 to 2 ft under the surface. At each sample site, the mesh bag and rope were replaced with clean, previously unused equipment.

## 2.2 SAMPLE PREPARATION

Laboratory procedures used to process and analyze the samples are discussed in detail in the QAPjP (Battelle, 1990b) and are summarized in this section. Detailed descriptions of the EPA methods used during this study are given in *Methods for Chemical Analysis of Water and Wastes* (EPA-600-1-79-020, March 1983). The procedures for processing the metals samples were the following.

- **Total recoverable metals.** The concentration of metals in an unfiltered sample following treatment with hot dilute mineral acid; described in Section 4.1.4 of EPA-600-1-79-020 (March 1983).

- **Dissolved metals.** Metals that will pass through a 0.45-$\mu$m-membrane filter in an unacidified sample; described in Section 4.1.1 of EPA-600-1-79-020 (March 1983).

- **Particulate metals** (also referred to as suspended metals). Metals that are retained by a 0.45-$\mu$m-membrane filter in an unacidified sample described in Section 4.1.2 of EPA-600-1-79-020 (March 1983).

Flow charts showing the processing steps that Battelle used for each of the metal phases measured in the this study are given in Battelle (1990b). Deviations from these EPA methods are given below.

### 2.2.1 Ambient Water Samples

Dissolved and particulate metals samples were processed in a Class 100 clean bench located within the wet laboratory of the OSV *Anderson*. Vacuum filtration at 5-10 psi was used to separate particulate from dissolved metals within about 2 h after sample collection. After the remaining sample had been discarded and the bottle rinsed with about 15 mL of filtrate, the metal-sample filtrate (dissolved phase) was placed back into its original sample container. Precombusted (350 °C) glass-fiber filters, with a 0.7-$\mu$m nominal pore size, were used for Hg filtrations, and Nuclepore membranes (0.4 $\mu$m) were used to filter samples for the other metals.

The same filter types were used to filter the acid-soluble phase approximately 2 days after acidification of the sample. Acid-soluble samples were processed in a Class 100 clean bench located within the wet laboratory of the OSV *Anderson*.

Particulate samples were stored frozen in acid-rinsed, airtight plastic petri dishes. Total-recoverable, acid-soluble, and dissolved-phase samples were acidified to pH 1.0 for mercury and at pH $<2.0$ for the other metals by using an ultrapure-grade nitric acid, and were stored at room temperature.

Samples for TSS were vacuum-filtered at 10 psi through preweighed 0.4-$\mu$m Nuclepore membrane filters. The total volume of water filtered was determined by using a graduated cylinder. Each filter was washed with three successive 10-mL rinses of deionized water (adjusted to pH 8 with ammonium hydroxide) to remove sea salt. Samples were stored in acid-rinsed, airtight, plastic petri dishes and air-dried in a Class 100 clean room prior to determination of the total mass of particles retained on the filter.

Duplicate PC and DOC sample aliquots from each sampling event were processed within about 2 h of sample collection. PC was collected on precombusted 25-mm glass-fiber filters, stored in an aluminum-foil pouch, and frozen until analysis. The DOC sample (filtrate) was sealed in a glass vial and also frozen until analysis. DOC and PC samples were shipped to Battelle's subcontractor, Chesapeake Bay Laboratory (CBL) in Solomons, Maryland, for analysis.

Samples for surface floc were processed in a Class 100 clean bench located within the wet laboratory of the OSV *Anderson*. Vacuum filtration at 5-10 psi was used to separate particulate from dissolved metals within about 6 h after sample collection. The particulate fraction was stored frozen in acid-rinsed, airtight, plastic petri dishes.

### 2.2.2 Tributary Samples

Dissolved and particulate metals samples were processed in a Class 100 clean room at Battelle in Duxbury, Massachusetts. Vacuum filtration at 5-10 psi was used to separate particulate from dissolved metals within about 1 day after sample collection. Precombusted (350 °C) glass-fiber filters (Whatman, GF/F), with a 0.7-$\mu$m nominal pore size, were used for Hg filtrations, and Nuclepore membranes (0.4 $\mu$m) were used to filter samples for the other metals.

Particulate samples were stored frozen in acid-rinsed, air-tight plastic petri dishes. Total-recoverable and dissolved phase samples were acidified to pH 1.0 for mercury and at pH $<2.0$ for the other metals by using an ultrapure-grade nitric acid.

Samples for TSS were vacuum-filtered at 10 psi through preweighed 0.4-$\mu$m Nuclepore membrane filters. The total volume of water filtered was determined by using a graduated cylinder. Each filter was washed with three successive 10-mL rinses of deionized water (adjusted to pH 8 with ammonium hydroxide) to remove sea salt. Samples were stored in acid-rinsed, airtight, plastic petri dishes and air-dried in a Class 100 clean room prior to determination of the total mass of particles retained on the filter.

Duplicate PC and DOC sample aliquots were processed from each municipal effluent within about 1 day of sample collection. PC was collected on precombusted, 25-mm glass-fibre filters stored in aluminum foil pouch, and frozen until analysis. The DOC sample (filtrate) was sealed in a glass vial and also frozen until analysis. Both DOC and PC samples were shipped to CBL for analysis.

### 2.2.3 Sample Digestion and Preconcentration

Both the IMSS samples and the tributary samples were prepared and analyzed using the same methods.

Total-recoverable metal analysis required additional sample preparation prior to analysis for all metals except Hg. A 100-mL aliquot of the sample was spiked with 500 $\mu$l $HNO_3$ and the sample was reduced in volume to about 20 mL by evaporation. Samples were diluted back to 100 mL and metals isolated by chelation/extraction (except As) before analysis. Arsenic samples were analyzed directly without preconcentration.

Ambient samples were concentrated by chelation and extraction at pH $< 1$ for Ag and at pH 4.5 for Cd, Cu, Ni, Pb, and Zn. Metals were chelated with ammonium-1-pyrrolidine dithiocarbamate/ diethylammonium diethyldithiocarbamate (APDC/DDDC) and extracted from the water with Freon (Danielson *et al.*, 1978). Prior to analysis, the solvent was acidified and the metals were backextracted into 2 mL of 10% nitric acid.

The particulate filter samples were digested in nitric acid (pH 1) at room temperature for about 2 days prior to analysis.

## 2.3 INSTRUMENTAL ANALYSIS

Instrumental analysis included either graphite furnace atomic absorption spectrometry (GFAAS), cold vapor atomic absorption spectrometry (CVAAS), or hydride atomic absorption spectrometry (HAAS). Samples were analyzed for Ag, Cd, Cu, Ni, Pb, and Zn by GFAAS after the chelation/extraction step for total-recoverable and dissolved metals.

For As analysis, a 5- to 10-mL sample aliquot was reacted with $NaBH_4$ to form a hydride prior to analysis by AAS.

For mercury analysis of ambient samples, about 250 mL of the acidified sample (total recoverable and dissolved) was processed by using a two-step analytical scheme described by Gill and Fitzgerald (1987) and Gill and Bruland (1990). The samples were reacted with $NaBH_4$ to cleave the C-Hg bonds, and the liberated Hg(II) was collected on a gilded-sand column and subsequently heated from the gold column for detection by CVAAS. Particulate sample digestates were analyzed directly by CVAAS without preconcentration.

## 2.4 SAMPLE CUSTODY

Field documentation consisted of log forms containing the project name, site number, sample-type designation, and an alphanumeric code. During field collection, log forms were completed and labels were affixed to the sample containers, thereby creating a link between the sample and data recorded on the log form. The log forms also contained the same alphanumeric code as their corresponding labels, ensuring the tracking of sample location and status.

The Chief Scientist recorded each sample on the Sample Transfer Record. Responsibility was relinquished by the Chief Scientist via signature at the time of receipt by laboratory personnel at Battelle. Transfer of samples to other laboratories took place at Battelle in Duxbury, Massachusetts.

Upon receipt of samples at Battelle, the Laboratory Sample Custodian examined the samples received, verified that the information recorded on the Sample Transfer Record was accurate, and logged the samples into the laboratory by signing the Sample Transfer Record. Date and time of receipt were also recorded. The sample identification numbers were used to track the samples through the laboratory.

Samples for PC/DOC analysis were shipped to CBL. The Battelle laboratory identification numbers were used to track the samples.

## 2.5 DATA QUALITY OBJECTIVES

A summary of the data-quality objectives for the analytical measurements is presented in Table 2. The reporting-limit goals (RLG) for metal analyses (Table 2) are one-fifth the most stringent WQC. Actual method detection limits (MDL) for metals analysis were determined from the instrument detection limits and the volume extracted. For all other analyses, historical MDLs are reported in Table 2 and were not redetermined for this project.

For metals analysis, accuracy was determined by using a standard reference material (SRM), and was also monitored via procedural blanks, equipment blanks, and matrix spike (MS) analysis. Precision was determined for metals via repeated analysis of an SRM and from analysis of laboratory duplicates of a sample and also field duplicates of a sample (samples were collected by splitting the sample stream so that the duplicate samples were collected simultaneously). For all other analyses, accuracy was determined from instrument calibration (salinity, TSS, PC) or analysis of matrix spikes (DOC only).

The quality-control results, which include procedural blanks, SRM recoveries, matrix spike recoveries, and analytical duplicate are listed by metal in Table 3. For this project, low-level procedural blanks were experienced for most of the analyses. Because the procedural blanks were low and consistent (see Appendix tables for procedural blanks), blank-corrected values are reported where appropriate for all metals except mercury. Blank corrections were determined and applied to each sample batch. All of the cadmium, copper, nickel, silver, and zinc and approximately 80% of the lead total recoverable results were blank-corrected. All of the silver, approximately 50% of the cadmium, copper, and lead, and 90% of the nickel and zinc dissolved results were blank-corrected. These corrections were generally less than 25% of the reported concentrations, except for total recoverable samples with extremely low concentrations (e.g., silver at station H8). Generally, the value of the Cd, Cu, Pb, and Zn blanks were <20%, <5%, <10%, and <30% of the lowest reported concentration, respectively. Blank corrections for Ni were higher, being approximately 50% of the reported concentration in the sample with the lowest Ni concentration.

The accuracy of the analytical results, as measured by the recovery of metals from standard reference materials and matrix spike recovery samples, was generally good, with most recoveries lying within ±15% of the expected values. Similarly, measures of analytical precision generally were within the data quality objectives of the program. Precision results of the IMSS field duplicates are listed in Table 4.

In a separate experiment, mercury was found to be stable for longer than the 30-day limit prescribed by EPA Methods for Chemical Analysis of Water and Wastes (EPA-600-1-79-020, March 1983). A mercury sample (station O4) collected in January 1991 under WA2-202 was reanalyzed in June 1991. The results differed by only 4% from the results of the first analysis conducted in February 1991.

## 3.0 RESULTS

The IMSS and the tributary sample analyses results for salinity, TSS, PC/DOC, and metals concentrations are listed in Table 3. Results from the IMSS sample analyses are plotted in Figure 2. Analyses of all dissolved As and acid-soluble Hg samples were not completed due to a programmatic decision to not apply the acid-soluble phase to the criteria being used in the TMDL/WLA and the need to complete the analysis of the hydrographic and turbidity data collected during the May IMSS survey (Battelle, 1991c).

Results from the surface floc samples collected during the IMSS are listed in Table 4.

## 4.0 CONCLUSIONS

This report presents a data set for the waters of NY/NJ Harbor that is of comparable quality to that reported for the January 1991 Ambient Evaluation Survey (Battelle, 1991b). Interpretation of the data presented in this report is beyond the scope of this project. However, the following are some observations and preliminary conclusions readily apparent from a cursory examination of the data.

- Surface and the bottom water total-recoverable mercury concentrations exceeded the EPA water quality criterion (0.025 $\mu$g/L) at stations H3, A3, and A7.

- Total-recoverable copper concentrations exceeded the EPA water quality criterion (2.0 $\mu$g/L) at all stations, times, and depths, with the exception of the surface sample from the station H8 (Hudson River). Because this is a freshwater station, the marine water quality criterion do not apply at this station.

- Dissolved copper concentrations do not exceed the EPA criterion, except at the stations located in the Arthur Kills region (A3 and A7)

- Bottom waters of station H3 exceeded the EPA WQC for lead and zinc in the total recoverable phase. All other lead and zinc values were below the EPA WQC.

- No exceedances of EPA WQC were observed for arsenic, cadmium, nickel, or silver.

- Bottom waters tended to have higher concentrations of metals than did surface waters.

- Except at station H3, variations in the total metals concentrations over the tidal cycle are small.

- Surface waters generally have lower TSS concentrations; the differences between surface and bottom waters are generally within a factor of 2 to 3.

- Variations in total suspended solids concentrations over the tidal cycle are small except at station H3.

- Changes in the total-recoverable metals concentrations over tidal cycles correspond closely with the changes in total suspended solids.

- Dissolved metals concentrations do not vary greatly over the tidal cycle. Changes are most likely related to variations in salinity (water type) than to remobilization of the metals from particulate matter.

- Particulate metal concentrations in the samples collected from near the sediment/water interface vary. The As, Hg, Cu, and Pb concentrations in the particulate samples from the Station H3 and H8 in the Hudson River are lower than at the other stations; Ni concentrations do not vary significantly.

- The particulate Hg results determined using room temperature pH 1 digestion are less than the difference between the total recoverable and dissolved measurement. This indicates particulate Hg is strongly bound to particles and released only with rigorous digestion techniques.

- The total-recoverable mercury concentration in the tributaries were less then the WQC. This implicates other sources as causes for high Hg in the Kills and at Hudson River station #3.

## 5.0 REFERENCES


Battelle. 1990a. Evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Work/quality assurance project plan submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1990b. Survey Plan for evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Final report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1991a. Initial Survey Report: Evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Final report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1991b. Results of the ambient and municipal sample interlaboratory comparison study in ambient waters and discharges to New York/New Jersey Harbor. Working draft report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1991c. Hydrographic results from the May 1992 Intensive Master Station Survey in New York/New Jersey Harbor. Letter report submitted to the Environmental Protection Agency under Work Assignment 2-213, Contract No. 68-C3-0105.

Danielson, L., B. Magnusson, and S. Westerlund. 1978. "An improved metal extraction procedure for the determination of trace metals in seawater by atomic absorption spectrometry with electrothermal atomization." Anal. Chem. Acta 98:47-57.

Gill, G.A., and W.F. Fitzgerald. 1987. "Picomolar mercury measurements in seawater and other materials using stannous chloride reduction and two-stage gold amalgamation with gas phase detection." Mar. Chem. 20:227-243.

Gill, G.A., and K.W. Bruland. 1990. "Mercury speciation in surface freshwater systems in California and other areas." Environ. Sci. Technol. 24:1392-1400.

Martin, T.D., J.W. O'Dell, and G.D. McKee. 1986. Evaluation of Method 200.1 determination of acid soluble metals. Draft report prepared by Inorganic Analyses Section, Physical and Chemical Methods Branch, Environmental Monitoring and Support Laboratory, Environmental Protection Agency Office of Research and Development, Cincinnati, OH.

Patterson, C.C., and D.M. Settle. 1976. "The reduction in orders of magnitude errors in lead analysis of biological materials and natural waters by evaluating and controlling the extent and sources of industrial lead contamination introduced during sample collecting and processing." In LaFleur, P.D. (Ed.), *Accuracy in Trace Analysis: Sampling, Sample Handling, Analysis.* National Bureau of Standards Spec. Pub. 422. U.S. Government Printing Office, Washington DC.

# Appendix

## FIGURES AND TABLES





Figure 1. Station Locations for May 1991 Intensive Master Station Survey.





Figure 2(a). Total Suspended Solids in New York/New Jersey Harbor Waters. .

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.





Figure 2(b). Particulate Carbon and Dissolved Organic Carbon in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.





**Figure 2(c).** **Arsenic Concentrations in New York/New Jersey Harbor Waters.**

**Data Are from May 1991 Intensive Master Station Survey.**
**E: Ebb. L: Low. F: Flood. H: High.**
**TR: Total Recoverable. D: Dissolved.**





**Figure 2(d).** Cadmium Concentrations in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved.





**Figure 2(e). Copper Concentrations in New York/New Jersey Harbor Waters.**

**Data Are from May 1991 Intensive Master Station Survey.**
**E: Ebb. L: Low. F: Flood. H: High.**
**TR: Total Recoverable. D: Dissolved.**





**Figure 2(f).** Lead Concentrations in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved.





Figure 2(g). Mercury Concentrations in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved. P: Particulate.





**Figure 2(h).** Nickel Concentrations in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved.





**Figure 2(i). Silver Concentrations in New York/New Jersey Harbor Waters.**

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved.





**Figure 2(j).** Zinc Concentrations in New York/New Jersey Harbor Waters.

Data Are from May 1991 Intensive Master Station Survey.
E: Ebb. L: Low. F: Flood. H: High.
TR: Total Recoverable. D: Dissolved.

Table 1. List of samples collected during the May 1991 Intensive Master Station Survey.[a]

| Time & Tidal Stage | Sample ID | Location | | Depth[b] (m) | Metals | | Mercury | | | | Split | Organic Part[c] | Surface Floc |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Lat. (N) | Long. (W) | | TR | D | TR | D | AS | P | | | |
| **May 15** | **Station 113** | | | | | | | | | | | | |
| 1055 H | NYJ00082 | 40°49.14′ | 73°58.17′ | 8.72 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1102 H | NYJ00083 | 40°49.14′ | 73°58.18′ | 2.87 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | — |
| 1348 E | NYJ00084 | 40°49.05′ | 73°58.23′ | 8.79 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | — |
| 1358 E | NYJ00085 | 40°49.05′ | 73°58.23′ | 2.70 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1406 E | NYJ00084SF | 40°49.05′ | 73°58.23′ | 9.30 | — | — | — | — | — | — | — | — | 1[d] |
| 1645 L | NYJ00086 | 40°49.03′ | 73°58.24′ | 7.35 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1652 L | NYJ00087 | 40°49.03′ | 73°58.24′ | 2.69 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1957 F | NYJ00088 | 40°49.09′ | 73°58.20′ | 8.17 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 2005 F | NYJ00089 | 40°49.09′ | 73°58.20′ | 3.09 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 2013 F | NYJ00088SF | 40°49.09′ | 73°58.20′ | 10.75 | — | — | — | — | — | — | — | — | 1 |
| **May 16** | **Station 118** | | | | | | | | | | | | |
| 1214 F | NYJ00090 | 41°32.33′ | 73°59.35′ | 10.18 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1220 F | NYJ00091 | 41°32.33′ | 73°59.35′ | 3.19 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1226 F | NYJ00090SF | 41°32.33′ | 73°59.34′ | 12.06 | — | — | — | — | — | — | — | — | 1[d] |
| 1507 H | NYJ00092 | 41°32.32′ | 73°59.33′ | 9.81 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1514 H | NYJ00093 | 41°32.32′ | 73°59.32′ | 3.19 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1819 E | NYJ00094 | 41°32.28′ | 73°59.30′ | 8.62 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | — |
| 1819 E | NYJ00098 | 41°32.28′ | 73°59.30′ | 8.62 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |

[a]Single samples for salinity and dissolved organic carbon as well as duplicate samples for particulate carbon and total suspended solids were collected whenever a metals total-recoverable sample was collected.
[b]Noncalibrated depth.
[c]Organic particulate.
[d]Sample collected but not filtered.
H: High; E: Ebb; L: Low; F: Flood.
TR: Total Recoverable; D: Dissolved; AS: Acid-soluble; P: Particulate.

Table 1. List of samples collected during the May 1991 Intensive Master Station Survey.[a] (continued)

| Time & Tidal Stage | Sample ID | Location | | Depth[b] (m) | Metals | | Mercury | | | | Split | Organic Part[c] | Surface Floc |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Lat. (N) | Long. (W) | | TR | D | TR | D | AS | P | | | |
| 1832 E | NYJ00095 | 41°32.24′ | 73°59.33′ | 2.94 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | — |
| 1842 E | NYJ00094SF | 41°32.25′ | 73°59.38′ | 11.14 | — | — | — | — | — | — | — | — | 1 |
| 2125 L | NYJ00096 | 41°32.24′ | 73°59.39′ | 9.58 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | — |
| 2136 L | NYJ00097 | 41°32.24′ | 73°59.38′ | 3.08 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | — |
| May 17 | Station B6 | | | | | | | | | | | | |
| 1155 H | NYJ00099 | 40°39.08′ | 74°03.45′ | 11.27 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1200 H | NYJ00100 | 40°39.20′ | 74°03.42′ | 4.06 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1449 E | NYJ00101 | 40°38.62′ | 74°03.08′ | 10.35 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 1 | — |
| 1449 E | NYJ00103 | 40°38.62′ | 74°03.08′ | 10.35 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1500 E | NYJ00102 | 40°38.62′ | 74°03.08′ | 3.51 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 1512 E | NYJ00101SF | 40°38.62′ | 74°03.06′ | 13.71 | — | — | — | — | — | — | — | — | 1[d] |
| 1739 L | NYJ00104 | 40°38.59′ | 74°03.13′ | 10.48 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 1746 L | NYJ00105 | 40°38.57′ | 74°03.13′ | 3.98 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 1 | — |
| 2049 F | NYJ00106 | 40°38.65′ | 74°03.07′ | 11.72 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 2056 F | NYJ00107 | 40°38.65′ | 74°03.07′ | 3.84 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 2104 F | NYJ00106SF | 40°38.64′ | 74°03.08′ | 13.94 | — | — | — | — | — | — | — | — | 1 |
| May 18 | Station A3 | | | | | | | | | | | | |
| 1016 F | NYJ00108 | 40°33.26′ | 74°14.68′ | 10.36 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1022 F | NYJ00109 | 40°33.30′ | 74°14.66′ | 3.20 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1030 F | NYJ00108SF | 40°33.27′ | 74°14.69′ | 11.64 | — | — | — | — | — | — | — | — | 1 |

[a]Single samples for salinity and dissolved organic carbon as well as duplicate samples for particulate carbon and total suspended solids were collected whenever a metals total-recoverable sample was collected.
[b]Noncalibrated depth.
[c]Organic particulate.
[d]Sample collected but not filtered.
H: High; E: Ebb; L: Low; F: Flood.
TR: Total Recoverable; D: Dissolved; AS: Acid-soluble; P: Particulate

Table 1. List of samples collected during the May 1991 Intensive Master Station Survey.[a] (continued)

| Time & Tidal Stage | Sample ID | Location | | Depth[b] (m) | Metals | | Mercury | | | | Split | Organic Part[c] | Surface Floc |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Lat. (N) | Long. (W) | | TR | D | TR | D | AS | P | | | |
| 1609 E | NYJ00112 | 40°33.29′ | 74°14.66′ | 8.70 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 1 | — |
| 1620 E | NYJ00113 | 40°33.28′ | 74°14.69′ | 2.32 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 1627 E | NYJ00112SF | 40°33.29′ | 74°14.69′ | 9.95 | — | — | — | — | — | — | — | — | 1[d] |
| 1845 L | NYJ00114 | 40°33.28′ | 74°14.68′ | 8.80 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1851 L | NYJ00115 | 40°33.28′ | 74°14.67′ | 3.18 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 1 | — |
| 1851 L | NYJ00116 | 40°33.28′ | 74°14.67′ | 3.18 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| May 19 | Station A7 | | | | | | | | | | | | |
| 0812 L | NYJ00117 | 40°40.21′ | 74°08.19′ | 9.59 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 0818 L | NYJ00118 | 40°40.22′ | 74°08.20′ | 3.11 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 1 | — |
| 1033 F | NYJ00119 | 40°40.25′ | 74°08.19′ | 12.03 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 1040 F | NYJ00120 | 40°40.26′ | 74°08.17′ | 3.55 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1046 F | NYJ00119SF | 40°40.26′ | 74°08.17′ | 12.68 | — | — | — | — | — | — | — | — | 1 |
| 1408 H | NYJ00121 | 40°40.36′ | 74°08.08′ | 11.32 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1414 H | NYJ00122 | 40°40.37′ | 74°08.06′ | 3.80 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 1710 E | NYJ00123 | 40°40.35′ | 74°08.07′ | 10.35 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 1 | — |
| 1710 E | NYJ00125 | 40°40.35′ | 74°08.07′ | 10.35 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1720 E | NYJ00124 | 40°40.37′ | 74°08.07′ | 3.53 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1727 E | NYJ00123SF | 40°40.37′ | 74°08.08′ | 12.96 | — | — | — | — | — | — | — | — | 1[d] |

[a]Single samples for salinity and dissolved organic carbon as well as duplicate samples for particulate carbon and total suspended solids were collected whenever a metals total-recoverable sample was collected.
[b]Noncalibrated depth.
[c]Organic particulate.
[d]Sample collected but not filtered.
H: High; E: Ebb; L: Low; F: Flood.
TR: Total Recoverable; D: Dissolved; AS: Acid-soluble; P: Particulate;

Table 1. List of samples collected during the May 1991 Intensive Master Station Survey.[a] (continued)

| Time & Tidal Stage | Sample ID | Location | | Depth[b] (m) | Metals | | Mercury | | | | Split | Organic Part[c] | Surface Floc |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Lat. (N) | Long. (W) | | TR | D | TR | D | AS | P | | | |
| May 20 | Station E4 | | | | | | | | | | | | |
| 0641 E | NYJ00126 | 40°47.79′ | 73°52.16′ | 10.49 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | — |
| 0649 E | NYJ00127 | 40°47.78′ | 73°52.16′ | 2.79 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 0655 E | NYJ00126SF | 40°47.79′ | 73°52.16′ | 13.40 | — | — | — | — | — | — | — | — | 1[d] |
| 1250 L | NYJ00128 | 40°47.79′ | 73°52.06′ | 8.86 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1256 L | NYJ00129 | 40°47.79′ | 73°52.06′ | 3.30 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 1 | — |
| 1327 F | NYJ00130 | 40°47.79′ | 73°52.06′ | 9.63 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 1 | — |
| 1327 F | NYJ00132 | 40°47.79′ | 73°52.06′ | 9.63 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1340 F | NYJ00131 | 40°47.79′ | 73°52.06′ | 3.13 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1346 F | NYJ00130SF | 40°47.80′ | 73°52.06′ | 11.98 | — | — | — | — | — | — | — | — | 2 |
| 1934 H | NYJ00133 | 40°47.79′ | 73°52.17′ | 10.91 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |
| 1948 H | NYJ00134 | 40°47.79′ | 73°52.18′ | 2.77 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | — |

[a]Single samples for salinity and dissolved organic carbon as well as duplicate samples for particulate carbon and total suspended solids were collected whenever a metals total-recoverable sample was collected.
[b]Noncalibrated depth.
[c]Organic particulate.
[d]Sample collected but not filtered.
H: High; E: Ebb; L: Low; F: Flood.
TR: Total Recoverable; D: Dissolved; AS: Acid-soluble; P: Particulate

Table 2. Data-Quality Objectives for Analytical Measurements.

| Parameter | Units | WQC[a] | RLG[b] | Accuracy[c] | Precision[d] | Method[e] |
|---|---|---|---|---|---|---|
| Ag | µg/L | 2.3 | 0.46 | ±20% | ±20% | Extraction/GFAAS |
| As | µg/L | 13 | 2.6 | ±20% | ±20% | HAAS |
| Cd | µg/L | 2.7 | 0.54 | ±20% | ±20% | Extraction/GFAAS |
| Cu | µg/L | 2.0 | 0.40 | ±20% | ±20% | Extraction/GFAAS |
| Hg | µg/L | 0.025 | 0.005 | ±20% | ±20% | Extraction/CVAAS |
| Ni | µg/L | 7.1 | 1.42 | ±20% | ±20% | Extraction/GFAAS |
| Pb | µg/L | 5.6 | 1.12 | ±20% | ±20% | Extraction/GFAAS |
| Zn | µg/L | 58 | 11.6 | ±20% | ±20% | Extraction/GFAAS |
| TSS | mg/L | NP[f] | 0.1 | ±10% | ±5% | Mass measurement |
| DOC | mg/L | NP | 0.24 | ±10% | ±5% | Digestion/infrared |
| PC | mg/L | NP | 0.063 | ±10% | ±10% | Elemental analyzer |

[a]Water quality criteria. Minimum marine chronic values are listed (from EPA or NYSDEC); except for Ag (acute value is listed. no chronic value is available).
[b]Reporting limit goals. The metal RLGs are one-fifth the corresponding minimum WQC as required for this program. Actual method detection limits (MDL) for the extraction/AAS methods are expected to be significantly lower than the RLGs. except for Ag. MDLs for direct-injection AAS methods are expected to be near or above the RLGs. Actual MDLs are given for the RLGs for total suspended solids (TSS), particulate carbon (PC), and dissolved organic carbon (DOC).
[c]Accuracy is determined as percent difference from true value (e.g., SRM, matrix spikes).
[d]Precision is determined as percent difference between replicate analysis.
[e]GFAAS: Graphite furnace atomic absorption spectrometry. HAAS: Hydride atomic absorption spectrometry. CVAAS: Cold vapor atomic absorption.
spectrometry.
[f]Not applicable.

Table 3(a). Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Salinity (‰). [a]

| Station | Tide | Surface[b] | Bottom[b] |
|---|---|---|---|
| E4 | Ebb | 23.83 | 24.18 |
| | Low | 23.75 | 23.89 |
| | Flood | 23.35 | 23.49 |
| | High | 23.98 | 24.46 |
| H3 | Ebb | 15.59 | 17.90 |
| | Low | 10.79 | 13.46 |
| | Flood | 9.93 | 11.43 |
| | High | 14.96 | 18.19 |
| H8 | Ebb | 0.10 | 0.10 |
| | Low | 0.09 | 0.10 |
| | Flood | 0.10 | 0.11 |
| | High | 0.10 | 0.11 |
| A7 | Ebb | 17.83 | 19.44 |
| | Low | 17.32 | 18.85 |
| | Flood | 17.98 | 19.58 |
| | High | 18.96 | 20.49 |
| A3 | Ebb | 21.61 | 22.86 |
| | Low | 20.08 | 21.61 |
| | Flood | 20.40 | 21.18 |
| B6 | Ebb | 22.87 | 23.13 |
| | Low | 22.42 | 25.76 |
| | Flood | 22.28 | 23.37 |
| | High | 24.22 | 26.17 |

[a]Salinity values not available for tributary stations.
[b]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively.

Table 3(b). Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Total Suspended Solids (mg/L).

| Station | Tide | Surface[a] | Bottom[a] |
|---|---|---|---|
| E4 | Ebb | 4.40 | 8.65 |
| | Low | 12.1 | 9.17 |
| | Flood | 7.97 | 9.88 |
| | High | 5.17 | 14.0 |
| H3 | Ebb | 15.9 | 169 |
| | Low | 132 | 409 |
| | Flood | 32.4 | 60.5 |
| | High | 17.9 | 317 |
| H8 | Ebb | 11.4 | 18.0 |
| | Low | 11.2 | 24.5 |
| | Flood | 6.79 | 7.27 |
| | High | 6.87 | 18.1 |
| A7 | Ebb | 9.50 | 7.58 |
| | Low | 13.8 | 14.1 |
| | Flood | 10.8 | 22.4 |
| | High | 8.99 | 12.5 |
| A3 | Ebb | 9.15 | 8.11 |
| | Low | 12.2 | 13.5 |
| | Flood | 10.7 | 25.5 |
| B6 | Ebb | 8.81 | 6.07 |
| | Low | 8.38 | 10.0 |
| | Flood | 8.78 | 8.75 |
| | High | 4.50 | 6.10 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]Not collected.
[c]Filter blank is based on 500-mL sample volume.

Table 3(b). Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Total Suspended Solids (mg/L). (continued)

| Station | Date | Surface[a] | Bottom[a] |
|---|---|---|---|
| H8 | 06/19/91 | 33.16 | 28.57 |
| | 06/23/91 | 16.20 | 14.31 |
| Hackensack | 06/04/91 | 13.96 | NC[b] |
| | 06/18/91 | 6.91 | NC |
| | 08/13/91 | 6.83 | NC |
| | 10/30/91 | 9.36 | NC |
| Passaic | 06/04/91 | 14.55 | NC |
| | 06/18/91 | 22.72 | NC |
| | 08/13/91 | 17.17 | NC |
| | 10/30/91 | 8.78 | NC |
| Raritan | 06/04/91 | 6.50 | NC |
| | 06/18/91 | 7.37 | NC |
| | 08/13/91 | 10.2 | NC |
| | 10/30/91 | 3.72 | NC |
| Quality Control Results | | | |
| Filter blank[c] | | | 0.33 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples which were collected <1 m below the surface.
[b]Not collected.
[c]Filter blank is based on 500-mL sample volume.

Table 3(c). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Particulate Carbon and Dissolved Organic Carbon (mg C/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | PC[b] | DOC[b] | PC | DOC |
| E4 | Ebb | 1.060 | 1.71 | 1.100 | 1.69 |
| | Low | 0.563 | 1.81 | 0.757 | 1.98 |
| | Flood | 0.657 | 1.77 | 0.666 | 1.78 |
| | High | 0.740 | 1.77 | 1.020 | 1.65 |
| H3 | Ebb | 0.906 | 2.19 | 9.120 | 2.36 |
| | Low | 4.760 | 2.63 | 17.100 | 2.44 |
| | Flood | 2.000 | 2.63 | 7.460 | 2.62 |
| | High | 0.696 | 2.20 | 13.600 | 2.33 |
| H8 | Ebb | 0.843 | 3.14 | 1.525 | 3.40 |
| | Low | 0.697 | 3.29 | 1.620 | 3.72 |
| | Flood | 0.747 | 3.43 | 1.060 | 3.47 |
| | High | 0.646 | 3.38 | 1.725 | 3.42 |
| A7 | Ebb | 1.420 | 2.67 | 0.666 | 2.18 |
| | Low | 1.090 | 2.65 | 1.310 | 2.24 |
| | Flood | 1.140 | 2.45 | 0.975 | 2.08 |
| | High | 1.160 | 2.26 | 0.639 | 2.16 |
| A3 | Ebb | 0.980 | 3.23 | 0.695 | 2.06 |
| | Low | 0.990 | 2.37 | 1.010 | 2.04 |
| | Flood | 0.945 | 2.56 | 1.350 | 2.19 |
| B6 | Ebb | 0.623 | 1.73 | 0.528 | 1.96 |
| | Low | 0.494 | 1.74 | 0.778 | 1.67 |
| | Flood | 0.557 | 1.83 | 0.499 | 2.00 |
| | High | 0.568 | 17.10 | 0.518 | 1.62 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected 1 m below the surface.
[b]PC: Particulate carbon. DOC: Dissolved organic carbon.
[c]Not collected.
[d]Not analyzed.

Table 3(c). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Particulate Carbon and Dissolved Organic Carbon (mg C/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | PC[b] | DOC[b] | PC | DOC |
| H8 | 06/19/91 | 0.924 | 3.17 | 0.360 | 4.11 |
| | 06/23/91 | 0.364 | 3.13 | 0.362 | 3.07 |
| Hackensack | 06/04/91 | 2.24 | 4.95 | NC[c] | NC |
| | 06/18/91 | 1.36 | 4.00 | NC | NC |
| | 08/13/91 | NA[d] | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |
| Passaic | 06/04/91 | 2.58 | 9.99 | NC | NC |
| | 06/18/91 | 2.93 | 6.61 | NC | NC |
| | 08/13/91 | NA | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |
| Raritan | 06/04/91 | 0.345 | 8.52 | NC | NC |
| | 06/18/91 | 0.735 | 51 | NC | NC |
| | 08/13/91 | NA | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |

| Quality Control Results | Particulate Carbon | Dissolved Organic Carbon |
|---|---|---|
| Matrix spike recovery (%)[e] | NP[f] | 104 ± 4 |
| Duplicates (RPD)[e] | 36 ±34 | 2 ± 1 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected 1 m below the surface.
[b]PC: Particulate carbon. DOC: Dissolved organic carbon.
[c]Not collected.
[d]Not analyzed.
[e]Mean ± standard deviation.
[f]Not applicable.

Table 3(d). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Arsenic (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 0.90 | NA[c] | 0.9 | 0.68 |
| | Low | 1.0 | NA | 1.1 | NA |
| | Flood | 1.0 | NA | 1.0 | NA |
| | High | 1.1 | NA | 1.1 | NA |
| H3 | Ebb | 1.2 | NA | 2.9 | NA |
| | Low | 1.9 | NA | 5.3 | NA |
| | Flood | 0.90 | NA | 2.0 | NA |
| | High | 1.1 | NA | 4.3 | NA |
| H8 | Ebb | 0.46 | NA | 0.56 | <0.19 |
| | Low | 0.43 | 0.18 | 0.51 | NA |
| | Flood | 0.43 | NA | 0.41 | NA |
| | High | 0.45 | NA | 0.56 | NA |
| A7 | Ebb | 1.2 | NA | 1.2 | NA |
| | Low | 1.5 | NA | 1.4 | NA |
| | Flood | 1.2 | NA | 1.5 | 0.78 |
| | High | 1.1 | 0.73 | 1.3 | NA |
| A3 | Ebb | 1.9 | 1.1 | 1.4 | 0.91 |
| | Low | 2.0 | NA | 2.0 | NA |
| | Flood | 1.7 | NA | 2.5 | NA |
| B6 | Ebb | 1.0 | 0.66 | 1.2 | 0.72 |
| | Low | 0.91 | 0.70 | 1.3 | 0.75 |
| | Flood | 1.2 | NA | 1.3 | NA |
| | High | 1.3 | NA | 1.3 | NA |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not analyzed.
[d]Not collected.
[e]Mean ± standard deviation or (range).
[f]Not applicable.

Table 3(d). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Arsenic (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 0.93 | NA[c] | 0.83 | NA |
| | 06/23/91 | 0.73 | NA | 0.73 | NA |
| Hackensack | 06/04/91 | 1.1 | NA | NC[d] | NC |
| | 06/18/91 | 0.68 | NA | NC | NC |
| | 08/13/91 | NA | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |
| Passaic | 06/04/91 | 1.0 | NA | NC | NC |
| | 06/18/91 | 1.5 | NA | NC | NC |
| | 08/13/91 | NA | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |
| Raritan | 06/04/91 | 1.4 | NA | NC | NC |
| | 06/18/91 | 1.2 | NA | NC | NC |
| | 08/13/91 | NA | NA | NC | NC |
| | 10/30/91 | NA | NA | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[e] | <0.03 (<0.03-0.05) | NP[f] |
| SRM recovery (%)[e] | 105 ± 3 | 70 (66-73) |
| Matrix spike recovery (%)[e] | 104 ± 13 | 116 |
| Duplicates (RPD)[e] | 2 ± 2 | NP |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected <1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not analyzed.
[d]Not collected.
[e]Mean ± standard deviation or (range).
[f]Not applicable.

Table 3(e). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Cadmium (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 0.068 | 0.070 | 0.070 | 0.051 |
| | Low | 0.069 | 0.064 | 0.076 | 0.070 |
| | Flood | 0.081 | 0.070 | 0.114 | 0.070 |
| | High | 0.066 | 0.065 | 0.079 | 0.060 |
| H3 | Ebb | 0.064 | 0.055 | 0.214 | 0.065 |
| | Low | 0.114 | 0.056 | 0.208 | 0.054 |
| | Flood | 0.067 | 0.048 | 0.109 | 0.057 |
| | High | 0.066 | 0.060 | 0.234 | 0.057 |
| H8 | Ebb | 0.024 | 0.013 | 0.064 | 0.008 |
| | Low | 0.022 | 0.010 | 0.045 | 0.011 |
| | Flood | 0.024 | 0.012 | 0.032 | 0.011 |
| | High | 0.028 | 0.015 | 0.061 | 0.011 |
| A7 | Ebb | 0.104 | 0.092 | 0.092 | 0.080 |
| | Low | 0.076 | 0.079 | 0.073 | 0.090 |
| | Flood | 0.100 | 0.082 | 0.115 | 0.063 |
| | High | 0.093 | 0.066 | 0.100 | 0.077 |
| A3 | Ebb | 0.134 | 0.086 | 0.100 | 0.060 |
| | Low | 0.148 | 0.129 | 0.126 | 0.099 |
| | Flood | 0.107 | 0.114 | 0.157 | 0.114 |
| B6 | Ebb | 0.075 | 0.050 | 0.078 | 0.050 |
| | Low | 0.063 | 0.047 | 0.068 | 0.034 |
| | Flood | 0.064 | 0.061 | 0.062 | 0.048 |
| | High | 0.040 | 0.054 | 0.020 | 0.044 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected <1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(e). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Cadmium (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 0.076 | 0.032 | 0.052 | 0.027 |
| | 06/23/91 | 0.036 | 0.038 | 0.032 | 0.027 |
| Hackensack | 06/04/91 | 0.039 | 0.012 | NC[c] | NC |
| | 06/18/91 | 0.028 | 0.022 | NC | NC |
| | 08/13/91 | 0.012 | 0.009 | NC | NC |
| | 10/30/91 | 0.016 | 0.014 | NC | NC |
| Passaic | 06/04/91 | 0.110 | 0.066 | NC | NC |
| | 06/18/91 | 0.212 | 0.090 | NC | NC |
| | 08/13/91 | 0.048 | 0.026 | NC | NC |
| | 10/30/91 | 0.149 | 0.153 | NC | NC |
| Raritan | 06/04/91 | 0.030 | 0.079 | NC | NC |
| | 06/18/91 | 0.054 | 0.043 | NC | NC |
| | 08/13/91 | 0.014 | 0.020 | NC | NC |
| | 10/30/91 | 0.035 | 0.034 | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[d,e] | 0.004 ± 0.003 | 0.002 ± 0.002 |
| SRM recovery (%)[e] | 92 ± 14 | 86 ± 22 |
| Matrix spike recovery (%)[e] | 83 ± 7 | 81 ± 7 |
| Duplicates (RPD)[e] | 4 ± 5 | 14 ± 11 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(f). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Copper (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 2.7 | 1.72 | 3.8 | 1.83 |
| | Low | 3.4 | 1.77 | 3.8 | 1.95 |
| | Flood | 3.8 | 1.83 | 4.4 | 1.82 |
| | High | 2.7 | 1.98 | 4.3 | 1.70 |
| H3 | Ebb | 3.2 | 1.71 | 23.8 | 1.72 |
| | Low | 12.2 | 1.94 | 36.6 | 1.72 |
| | Flood | 4.1 | 1.98 | 10.2 | 1.95 |
| | High | 3.1 | 1.75 | 29.2 | 1.63 |
| H8 | Ebb | 1.9 | 1.50 | 3.2 | 1.44 |
| | Low | 1.8 | 1.40 | 2.6 | 1.38 |
| | Flood | 1.6 | 1.46 | 2.3 | 1.25 |
| | High | 1.9 | 1.53 | 3.7 | 1.53 |
| A7 | Ebb | 3.8 | 2.22 | 3.5 | 1.88 |
| | Low | 3.9 | 2.04 | 4.3 | 1.96 |
| | Flood | 3.5 | 1.86 | 6.2 | 1.96 |
| | High | 3.5 | 2.01 | 4.1 | 1.72 |
| A3 | Ebb | 6.4 | 2.36 | 5.0 | 2.05 |
| | Low | 6.4 | 2.39 | 6.5 | 2.13 |
| | Flood | 5.3 | 2.50 | 12.4 | 2.41 |
| B6 | Ebb | 3.3 | 1.56 | 2.7 | 1.53 |
| | Low | 2.9 | 1.47 | 3.2 | 1.13 |
| | Flood | 2.8 | 1.79 | 2.9 | 1.37 |
| | High | 1.7 | 1.56 | 2.6 | 1.30 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(f). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Copper (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 4.72 | 2.03 | 3.59 | 1.90 |
| | 06/23/91 | 3.16 | 2.13 | 3.39 | 2.64 |
| Hackensack | 06/04/91 | 18.09 | 9.85 | NC[c] | NC |
| | 06/18/91 | 16.32 | 10.97 | NC | NC |
| | 08/13/91 | 5.44 | 3.68 | NC | NC |
| | 10/30/91 | 6.16 | 6.21 | NC | NC |
| Passaic | 06/04/91 | 6.56 | 3.94 | NC | NC |
| | 06/18/91 | 12.58 | 6.96 | NC | NC |
| | 08/13/91 | 5.55 | 3.06 | NC | NC |
| | 10/30/91 | 4.97 | 4.59 | NC | NC |
| Raritan | 06/04/91 | 3.03 | 2.61 | NC | NC |
| | 06/18/91 | 4.04 | 2.98 | NC | NC |
| | 08/13/91 | 3.65 | 2.73 | NC | NC |
| | 10/30/91 | 4.24 | 4.00 | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[d,e] | 0.09 ± 0.06 | 0.03 ± 0.05 |
| SRM recovery (%)[e] | 99 ± 8 | 100 ± 7 |
| Matrix spike recovery (%)[e] | 96 ± 7 | 100 ± 2 |
| Duplicates (RPD)[e] | 3 ± 3 | 5 ± 8 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(g). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Lead (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 1.04 | 0.176 | 2.02 | 0.169 |
| | Low | 1.57 | 0.198 | 2.27 | 0.213 |
| | Flood | 1.80 | 0.247 | 2.32 | 0.237 |
| | High | 1.01 | 0.170 | 2.39 | 0.158 |
| H3 | Ebb | 2.09 | 0.126 | 22.2 | 0.206 |
| | Low | 2.27 | 0.092 | 29.6 | 0.108 |
| | Flood | 3.49 | 0.100 | 2.07 | 0.130 |
| | High | 2.15 | 0.125 | 12.9 | 0.186 |
| H8 | Ebb | 0.91 | 0.075 | 2.25 | 0.095 |
| | Low | 0.81 | 0.080 | 1.94 | 0.087 |
| | Flood | 0.61 | 0.073 | 1.36 | 0.064 |
| | High | 1.15 | 0.077 | 2.86 | 0.090 |
| A7 | Ebb | 2.54 | 0.165 | 2.14 | 0.181 |
| | Low | 2.72 | 0.190 | 3.00 | 0.192 |
| | Flood | 2.43 | 0.172 | 4.54 | 0.206 |
| | High | 2.10 | 0.192 | 2.64 | 0.192 |
| A3 | Ebb | 3.33 | 0.225 | 2.52 | 0.159 |
| | Low | 3.61 | 0.280 | 3.55 | 0.217 |
| | Flood | 2.92 | 0.286 | 1.54 | 0.117 |
| B6 | Ebb | 2.07 | 0.206 | 1.52 | 0.186 |
| | Low | 1.44 | 0.165 | 2.34 | 0.154 |
| | Flood | 1.36 | 0.225 | 1.60 | 0.175 |
| | High | 0.74 | 0.175 | 1.49 | 0.202 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(g). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Lead (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 3.14 | 0.147 | 2.17 | 0.151 |
| | 06/23/91 | 1.45 | 0.091 | 1.33 | 0.094 |
| Hackensack | 06/04/91 | 1.37 | 0.068 | NC[c] | NC |
| | 06/18/91 | 3.58 | 1.311 | NC | NC |
| | 08/13/91 | 0.28 | 0.038 | NC | NC |
| | 10/30/91 | 0.92 | 0.87 | NC | NC |
| Passaic | 06/04/91 | 4.63 | 0.474 | NC | NC |
| | 06/18/91 | 14.72 | 1.931 | NC | NC |
| | 08/13/91 | 4.68 | 0.523 | NC | NC |
| | 10/30/91 | 3.08 | 3.10 | NC | NC |
| Raritan | 06/04/91 | 0.55 | 0.256 | NC | NC |
| | 06/18/91 | 0.66 | 0.181 | NC | NC |
| | 08/13/91 | 0.77 | 0.212 | NC | NC |
| | 10/30/91 | 0.60 | 0.28 | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[d,e] | 0.01 ± 0.02 | 0.007 ± 0.007 |
| SRM recovery (%)[e] | 89 ± 37 | 70 ± 21 |
| Matrix spike recovery (%)[e] | 97 ± 8 | 96 ± 4 |
| Duplicates (RPD)[e] | 4 ± 2 | 5 ± 5 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(h). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Mercury (μg/L).

| Station | Tide | Surface[a] | | | Bottom[a] | | |
|---|---|---|---|---|---|---|---|
| | | TR[b] | D[b] | P[b] | TR | D | P |
| E4 | Ebb | 0.0072 | 0.0027 | NC[c] | 0.0109 | <0.0028 | NC |
| | Low | 0.0118 | <0.0028 | NC | 0.0150 | <0.0028 | NC |
| | Flood | 0.0131 | <0.0028 | NC | 0.0161 | 0.0030 | NC |
| | High | 0.0063 | <0.0028 | NC | 0.0138 | <0.0028 | NC |
| H3 | Ebb | 0.0195 | 0.0030 | <0.00023 | 0.1818 | <0.0028 | <0.00024 |
| | Low | 0.0621 | <0.0028 | 0.00038 | 0.2539 | 0.0031 | <0.00028 |
| | Flood | 0.0221 | 0.0033 | <0.00022 | 0.0127 | <0.0028 | <0.00023 |
| | High | 0.0186 | <0.0028 | <0.00022 | 0.1873 | 0.0032 | <0.00026 |
| H8 | Ebb | 0.0032 | <0.0028 | <0.00023 | 0.0077 | 0.0035 | <0.00023 |
| | Low | 0.0051 | 0.0048 | <0.00022 | 0.0048 | <0.0028 | <0.00025 |
| | Flood | <0.0020 | 0.0029 | <0.00022 | 0.0071 | 0.0027 | <0.00024 |
| | High | 0.0021 | 0.0027 | <0.00022 | 0.0060 | 0.0038 | <0.00023 |
| A7 | Ebb | 0.0321 | 0.0037 | NC | 0.0266 | <0.0028 | NC |
| | Low | 0.0513 | <0.0028 | NC | 0.0598 | 0.0040 | NC |
| | Flood | 0.0282 | 0.0041 | NC | 0.0609 | <0.0028 | NC |
| | High | 0.0190 | <0.0028 | NC | 0.0351 | <0.0028 | NC |
| A3 | Ebb | 0.0433 | 0.0036 | NC | 0.0315 | <0.0028 | NC |
| | Low | 0.0527 | 0.0024 | NC | 0.0562 | <0.0028 | NC |
| | Flood | 0.0409 | 0.0034 | NC | 0.0416 | <0.0028 | NC |
| B6 | Ebb | 0.0202 | <0.0028 | NC | 0.0143 | <0.0028 | NC |
| | Low | 0.0125 | <0.0028 | NC | 0.0263 | <0.0028 | NC |
| | Flood | 0.0098 | <0.0028 | NC | 0.0166 | <0.0028 | NC |
| | High | 0.0085 | <0.0028 | NC | 0.0140 | <0.0028 | NC |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved. P: Particulate.
[c]Not collected.
[d]Procedural blank is based on 250-mL sample volume.
[e]Mean ± standard deviation.

Table 3(h). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Mercury (µg/L). (continued)

| Station | Date | Surface[a] | | | Bottom[a] | | |
|---|---|---|---|---|---|---|---|
| | | TR[b] | D[b] | P[b] | TR | D | P |
| H8 | 06/19/91 | 0.0044 | 0.0045 | 0.00033 | 0.0060 | 0.0048 | 0.00029 |
| | 06/23/91 | 0.0042 | 0.0081 | 0.00017 | 0.0074 | 0.0038 | <0.00018 |
| Hackensack | 06/04/91 | 0.0065 | 0.0086 | 0.00055 | NC[c] | NC | NC |
| | 06/18/91 | 0.0028 | 0.0062 | 0.00038 | NC | NC | NC |
| | 08/13/91 | 0.0039 | 0.0054 | NR[d] | NC | NC | NC |
| | 10/30/91 | 0.0024 | 0.0968 | NR | NC | NC | NC |
| Passaic | 06/04/91 | 0.0138 | 0.0063 | 0.00038 | NC | NC | NC |
| | 06/18/91 | 0.0183 | 0.0103 | <0.00017 | NC | NC | NC |
| | 08/13/91 | 0.0085 | 0.0052 | NR | NC | NC | NC |
| | 10/30/91 | 0.0043 | 0.0080 | NR | NC | NC | NC |
| Raritan | 06/04/91 | 0.0054 | 0.0042 | <0.00018 | NC | NC | NC |
| | 06/18/91 | 0.0049 | 0.0115 | <0.00017 | NC | NC | NC |
| | 08/13/91 | 0.0032 | 0.0056 | NR | NC | NC | NC |
| | 10/30/91 | 0.0043 | 0.0140 | NR | NC | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural Blank (µg/L)[e,f] | 0.002 ± 0.002 | 0.002 ± 0.001 |
| Blank Spike Recovery (%)[f] | 99 ± 13 | 95 ± 8 |
| Matrix Spike Recovery (%)[f] | 78 ± 25 | 90 ± 11 |
| Duplicates (%RPD)[f] | 20 ± 9 | 6 ± 8 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved. P: Particulate.
[c]Not collected.
[d]Not reported.
[e]Procedural blank is based on 250-mL sample volume.
[f]Mean ± standard deviation.

Table 3(i). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Nickel (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 1.4 | 1.44 | 1.8 | 1.27 |
| | Low | 1.7 | 1.46 | 1.4 | 1.67 |
| | Flood | 2.0 | 1.57 | 2.4 | 1.53 |
| | High | 1.6 | 1.52 | 1.8 | 1.37 |
| H3 | Ebb | 2.0 | 0.91 | 6.3 | 0.91 |
| | Low | 4.3 | 0.68 | 9.3 | 0.74 |
| | Flood | 1.9 | 0.70 | 3.9 | 0.78 |
| | High | 1.7 | 0.89 | 7.8 | 0.93 |
| H8 | Ebb | 1.0 | 0.38 | 1.9 | 0.40 |
| | Low | 0.7 | 0.46 | 1.4 | 0.34 |
| | Flood | 0.7 | 0.42 | 1.0 | 0.34 |
| | High | 1.2 | 0.35 | 1.6 | 0.36 |
| A7 | Ebb | 2.2 | 2.06 | 2.0 | 1.79 |
| | Low | 1.5 | 2.26 | 1.2 | 1.89 |
| | Flood | 2.1 | 1.88 | 3.2 | 1.54 |
| | High | 2.1 | 1.65 | 1.5 | 1.61 |
| A3 | Ebb | 2.7 | 1.90 | 2.1 | 1.53 |
| | Low | 2.7 | 2.74 | 2.4 | 2.14 |
| | Flood | 1.7 | 2.75 | 3.7 | 1.13 |
| B6 | Ebb | 1.8 | 1.26 | 1.8 | 1.18 |
| | Low | 1.5 | 1.08 | 1.6 | 0.90 |
| | Flood | 1.6 | 1.18 | 1.4 | 0.94 |
| | High | 1.3 | 1.04 | 1.2 | 0.83 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(i). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Nickel (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 1.58 | 0.55 | 1.17 | 0.50 |
| | 06/23/91 | 0.92 | 0.71 | 1.31 | 0.60 |
| Hackensack | 06/04/91 | 0.91 | 0.85 | NC[c] | NC |
| | 06/18/91 | 0.59 | 0.83 | NC | NC |
| | 08/13/91 | 0.39 | 0.43 | NC | NC |
| | 10/30/91 | 0.72 | 0.69 | NC | NC |
| Passaic | 06/04/91 | 3.12 | 3.06 | NC | NC |
| | 06/18/91 | 4.20 | 3.17 | NC | NC |
| | 08/13/91 | 1.92 | 1.39 | NC | NC |
| | 10/30/91 | 3.80 | 3.42 | NC | NC |
| Raritan | 06/04/91 | 1.14 | 1.74 | NC | NC |
| | 06/18/91 | 1.28 | 1.75 | NC | NC |
| | 08/13/91 | 1.30 | 1.03 | NC | NC |
| | 10/30/91 | 1.56 | 1.57 | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[d,e] | 0.4 ± 0.2 | 0.2 ± 0.1 |
| SRM recovery (%)[f] | 97 ± 22 | 92 ± 7 |
| Matrix spike recovery (%)[g] | 100 ± 11 | 94 ± 7 |
| Duplicates (RPD)[g] | 9 ± 5 | 5 ± 7 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected <1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(j). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Silver (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 0.047 | 0.0102 | 0.054 | 0.0087 |
| | Low | 0.067 | 0.0063 | 0.085 | 0.0091 |
| | Flood | 0.067 | 0.0074 | 0.080 | 0.0076 |
| | High | 0.037 | 0.0093 | 0.084 | 0.0056 |
| H3 | Ebb | 0.087 | 0.0166 | 0.218 | 0.0197 |
| | Low | 0.310 | 0.0111 | 0.352 | 0.0083 |
| | Flood | 0.129 | 0.0060 | 0.301 | 0.0102 |
| | High | 0.090 | 0.0078 | 0.337 | 0.0104 |
| H8 | Ebb | 0.006 | 0.0021 | 0.026 | 0.0015 |
| | Low | 0.008 | 0.0018 | 0.011 | 0.0101 |
| | Flood | 0.007 | 0.0008 | 0.008 | 0.0010 |
| | High | 0.005 | 0.0019 | 0.009 | 0.0004 |
| A7 | Ebb | 0.048 | 0.0048 | 0.052 | 0.0052 |
| | Low | 0.056 | 0.0020 | 0.070 | 0.0087 |
| | Flood | 0.048 | 0.0034 | 0.083 | 0.0045 |
| | High | 0.036 | 0.0046 | 0.081 | 0.0069 |
| A3 | Ebb | 0.067 | 0.0053 | 0.041 | 0.0090 |
| | Low | 0.084 | 0.0059 | 0.083 | 0.0057 |
| | Flood | 0.064 | 0.0063 | 0.163 | 0.0106 |
| B6 | Ebb | 0.079 | 0.0098 | 0.062 | 0.0100 |
| | Low | 0.052 | 0.0091 | 0.064 | 0.0054 |
| | Flood | 0.070 | 0.0100 | 0.088 | 0.0079 |
| | High | 0.025 | 0.0092 | 0.069 | 0.0103 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not detected above the blank.
[d]Not collected.
[e]Procedural blank is based on 100-mL sample volume.
[f]Mean ± standard deviation.
[g]Not applicable.

Table 3(j). Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Silver (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 0.023 | 0.005 | 0.017 | 0.008 |
| | 06/23/91 | 0.006 | 0.007 | 0.005 | 0.002 |
| Hackensack | 06/04/91 | ND[c] | 0.005 | NC[d] | NC |
| | 06/18/91 | ND | 0.001 | NC | NC |
| | 08/13/91 | 0.004 | ND | NC | NC |
| | 10/30/91 | ND[e] | 0.004[e] | NC | NC |
| Passaic | 06/04/91 | 0.089 | 0.018 | NC | NC |
| | 06/18/91 | 0.180 | 0.090 | NC | NC |
| | 08/13/91 | 0.060 | ND | NC | NC |
| | 10/30/91 | 0.054[e] | 0.014[e] | NC | NC |
| Raritan | 06/04/91 | ND | 0.010 | NC | NC |
| | 06/18/91 | ND | 0.010 | NC | NC |
| | 08/13/91 | 0.021 | ND | NC | NC |
| | 10/30/91 | 0.010[e] | 0.007[e] | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[f,g] | 0.01 ± 0.007 | 0.01 ± 0.006 |
| SRM recovery (%)[g] | NP[h] | NP |
| Matrix spike recovery (%)[g] | 105 ± 48 | 64 ± 28 |
| Duplicates (RPD)[g] | 32 ± 49 | 30 ± 25 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not detected above the blank.
[d]Not collected.
[e]Inefficient silver extraction.
[f]Procedural blank is based on 100-mL sample volume.
[g]Mean ± standard deviation.
[h]Not applicable.

Table 3(k). Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Zinc (μg/L).

| Station | Tide | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| E4 | Ebb | 6.3 | 5.20 | 6.6 | 5.16 |
| | Low | 8.6 | 5.09 | 6.8 | 5.35 |
| | Flood | 7.9 | 5.68 | 8.5 | 5.44 |
| | High | 6.5 | 5.32 | 8.2 | 4.49 |
| H3 | Ebb | 7.7 | 3.35 | 50.1 | 3.57 |
| | Low | 27.3 | 1.93 | 74.4 | 2.69 |
| | Flood | 8.5 | 1.70 | 10.0 | 2.58 |
| | High | 7.6 | 3.03 | 64.8 | 3.27 |
| H8 | Ebb | 2.8 | 0.52 | 6.4 | 0.66 |
| | Low | 1.9 | 0.63 | 6.6 | 0.73 |
| | Flood | 2.3 | 0.56 | 4.2 | 0.84 |
| | High | 3.3 | 0.66 | 8.5 | 0.57 |
| A7 | Ebb | 10.5 | 7.92 | 9.0 | 7.86 |
| | Low | 7.4 | 7.86 | 6.2 | 7.81 |
| | Flood | 9.3 | 7.60 | 12.2 | 7.68 |
| | High | 8.9 | 7.78 | 6.5 | 6.84 |
| A3 | Ebb | 11.0 | 7.22 | 7.1 | 5.08 |
| | Low | 12.8 | 10.40 | 10.2 | 7.23 |
| | Flood | 6.6 | 9.38 | 19.6 | 7.95 |
| B6 | Ebb | 6.4 | 4.57 | 5.7 | 4.38 |
| | Low | 5.2 | 4.00 | 5.5 | 2.85 |
| | Flood | 6.5 | 4.75 | 5.9 | 3.43 |
| | High | 4.3 | 3.88 | 5.5 | 2.89 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 3(k). Results from the New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Zinc (μg/L). (continued)

| Station | Date | Surface[a] | | Bottom[a] | |
|---|---|---|---|---|---|
| | | TR[b] | D[b] | TR | D |
| H8 | 06/19/91 | 8.50 | 3.20 | 6.25 | 2.52 |
| | 06/23/91 | 3.57 | 2.48 | 4.11 | 3.46 |
| Hackensack | 06/04/91 | 7.84 | 1.59 | NC[c] | NC |
| | 06/18/91 | 4.05 | 1.85 | NC | NC |
| | 08/13/91 | 1.54 | 2.26 | NC | NC |
| | 10/30/91 | 2.42 | 2.09 | NC | NC |
| Passaic | 06/04/91 | 14.78 | 6.49 | NC | NC |
| | 06/18/91 | 32.06 | 16.96 | NC | NC |
| | 08/13/91 | 8.05 | 3.81 | NC | NC |
| | 10/30/91 | 10.09 | 10.10 | NC | NC |
| Raritan | 06/04/91 | 3.91 | 3.73 | NC | NC |
| | 06/18/91 | 7.03 | 6.44 | NC | NC |
| | 08/13/91 | 4.15 | 4.83 | NC | NC |
| | 10/30/91 | 7.30 | 7.76 | NC | NC |

| Quality Control Results | Total Recoverable | Dissolved |
|---|---|---|
| Procedural blank (μg/L)[d,e] | 0.4 ± 0.2 | 0.2 ± 0.1 |
| SRM recovery (%)[e] | 105 ± 25 | 109 ± 13 |
| Matrix spike recovery (%)[e] | 95 ± 23 | 101 ± 7 |
| Duplicates (RPD)[e] | 16 ± 14 | 6 ± 4 |

[a]Samples were collected at the surface and bottom, at 1/4 and 3/4 depth in the water column, respectively, except for the New Jersey Tributary samples, which were collected < 1 m below the surface.
[b]TR: Total recoverable. D: Dissolved.
[c]Not collected.
[d]Procedural blank is based on 100-mL sample volume.
[e]Mean ± standard deviation.

Table 4. Precision Results of the Field-duplicate Samples Collected for the New York/New Jersey Harbor Intensive Master Station Survey — Relative Percent Difference.

| Station | As | Cd | Cu | Pb | Hg | Ni | Ag | Zn |
|---|---|---|---|---|---|---|---|---|
| **Total Recoverable** | | | | | | | | |
| E4 | 10 | 64 | 2 | <1 | 14 | 33 | 9 | 12 |
| H8 | 21 | 9 | 16 | 14 | 54 | 8 | 65 | 35 |
| A7 | 8 | 2 | 6 | 2 | 12 | <1 | 2 | 33 |
| A3 | 15 | 12 | 6 | 6 | 4 | 15 | 16 | 38 |
| B6 | 9 | 25 | 2 | 3 | 5 | 20 | 44 | 18 |
| **Dissolved** | | | | | | | | |
| E4 | NA[a] | 7 | 3 | 3 | ND[b] | 7 | 11 | 2 |
| H8 | ND | 25 | 7 | 23 | 5 | 2 | 120 | 21 |
| A7 | NA | 10 | 2 | 3 | ND | 2 | 29 | 4 |
| A3 | NA | 2 | 11 | 8 | ND | 3 | 17 | 1 |
| B6 | 1 | 38 | 1 | 2 | ND | 1 | 12 | 8 |

[a]Not analyzed.
[b]Not detected. At least one of the field duplicates was below the detection limit.

Table 5. Results from New York/New Jersey Harbor Intensive Master Station and Tributary Surveys — Surface Floc (μg/g particulate matter).

| Station | Ag | As | Cd | Cu | Hg | Ni | Pb | Zn |
|---|---|---|---|---|---|---|---|---|
| E4 | <0.162 | 11.54 | 1.18 | 185.2 | <0.0651 | 12.5 | 184.0 | 188 |
| H3 | <0.006 | 0.76 | 0.90 | 91.8 | <0.0025 | 15.6 | 77.3 | 205 |
| H8 | <0.012 | 0.72 | 1.56 | 39.2 | <0.0048 | 12.7 | 50.4 | 129 |
| A7 | <0.049 | 11.18 | 1.76 | 141.9 | <0.0197 | 12.3 | 179.2 | 237 |
| A3 | <0.092 | 24.69 | 2.38 | 276.1 | <0.0366 | 18.4 | 238.9 | 386 |
| B6 | <0.065 | 11.47 | 1.17 | 122.5 | <0.0258 | 12.8 | 131.2 | 195 |
| Quality Control | | | | | | | | |
| Filter blank[a] | <0.024 | <0.39 | 0.012 | <0.53 | NR[b] | <0.38 | <0.08 | 0.26 |

[a]Filter blanks based on a 0.017-g mean particulate mass.
[b]Not reported; contamination suspected.

Battelle
Ocean Sciences
397 Washington Street
Duxbury, Massachusetts 02332
Telephone. (617) 934-0571

January 20, 1992

Mr. Seth Ausubel
EPA Region II
26 Federal Plaza
New York, NY 10278

Dear Mr. Ausubel:

Enclosed are corrections and additions to the final data report *Evaluation of Trace-Metal Levels in Ambient Waters and Tributaries to New York/New Jersey Harbor for Waste Load Allocation*. Please replace page 8 you currently have with the one accompanying this letter. We have not changed the water quality criteria for Cu, Pb, and Zn elsewhere in the document because these must remain consistent with the approved Quality Assurance Project Plan.

If you have questions regarding this report please call Dr. Carlton Hunt or Dr. Damian Shea at (617) 934-0571.

Sincerely,

Damian Shea

Damian Shea, Ph.D.

Enclosure

## 4.0 CONCLUSIONS

This report presents a data set for the waters of NY/NJ Harbor that is of comparable quality to that reported for the January 1991 Ambient Evaluation Survey (Battelle, 1991b). Interpretation of the data presented in this report is beyond the scope of this project. However, the following are some observations and preliminary conclusions readily apparent from a cursory examination of the data.

- Surface and the bottom water total-recoverable mercury concentrations exceeded the EPA water quality criterion (0.025 µg/L) at stations H3, A3, and A7.

- Total-recoverable copper concentrations exceed the EPA water quality criterion (2.9 µg/L) at stations A3, A7, E4, and H3 at all times and depths, with the exception of the surface ebb and high tide from station E4. There were two exceedences for both stations H8 and B6. Station H8 exceeded WQC during ebb and high tide at the bottom and station B6 during ebb tide at the surface and during low tide at the bottom. Because station H8 is a freshwater station, the marine water quality criterion do not apply at this station.

- Dissolved copper concentrations do not exceed the EPA criterion.

- Bottom waters of station H3 exceeded EPA WQC for lead, nickel, and zinc during low tide, lead and nickel during high tide, and lead during ebb tide. All other lead, nickel, and zinc values were below EPA WQC.

- No exceedances of EPA WQC were observed for arsenic, cadmium, or silver.

- Bottom waters tended to have higher concentrations of metals than did surface waters.

- Except at station H3, variations in the total metals concentrations over the tidal cycle are small.

- Surface waters generally have lower TSS concentrations; the differences between surface and bottom waters are generally within a factor of 2 to 3.

- Variations in total suspended solids concentrations over the tidal cycle are small except at station H3.

- Changes in the total-recoverable metals concentrations over tidal cycles correspond closely with the changes in total suspended solids.

- Dissolved metals concentrations do not vary greatly over the tidal cycle. Changes are most likely related to variations in salinity (water type) than to remobilization of the metals from particulate matter.

- Particulate metal concentrations in the samples collected from near the sediment/water interface vary. The As, Hg, Cu, and Pb concentrations in the particulate samples from the Station H3 and H8 in the Hudson River are lower than at the other stations; Ni concentrations do not vary significantly.

- The particulate Hg results determined using room temperature pH 1 digestion are less than the difference between the total recoverable and dissolved measurement. This indicates particulate Hg is strongly bound to particles and released only with rigorous digestion techniques.

- The total-recoverable mercury concentration in the tributaries were less then the WQC. This implicates other sources as causes for high Hg in the Kills and at Hudson River station #3.

## 5.0 REFERENCES


Battelle. 1990a. Evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Work/quality assurance project plan submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1990b. Survey Plan for evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Final report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1991a. Initial Survey Report: Evaluation of trace-metal levels in ambient waters and discharges to New York/New Jersey Harbor for waste load allocation. Final report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.

Battelle. 1991b. Results of the ambient and municipal sample interlaboratory comparison study in ambient waters and discharges to New York/New Jersey Harbor. Working draft report submitted to the Environmental Protection Agency under Work Assignment 2-202, Contract No. 68-C3-0105.